



# ENHANCING TRAUMA CARE IN INPATIENT PSYCHIATRY: THE ROLE OF MENTAL HEALTH NURSES IN IMPLEMENTING TRAUMA-FOCUSED COGNITIVE BEHAVIORAL THERAPY (TF-CBT)

Santhosh Kumar. J<sup>1</sup>

<sup>1</sup>Associate Professor, Department of Mental Health (Psychiatric) Nursing,  
Amrita College of Nursing, Amrita Vishwa Vidyapeetham, Kochi, Kerala, India.

## ABSTRACT

Trauma is a pervasive issue within inpatient psychiatric care, where a significant number of patients have trauma histories that contribute to their mental health disorders. Trauma-Focused Cognitive Behavioral Therapy (TF-CBT) has emerged as an effective, evidence-based treatment for trauma-related mental health conditions, particularly for patients suffering from post-traumatic stress disorder (PTSD). Mental health nurses, as front-line caregivers, have a critical role in the application of TF-CBT, offering trauma-informed care that can significantly enhance patient outcomes. This article provides a comprehensive review of how mental health nurses can implement TF-CBT in inpatient psychiatric settings, discusses the competencies required, addresses barriers to implementation, and offers recommendations for practice. Through this review, we emphasize the potential for improved patient care and outcomes when trauma is adequately addressed in psychiatric care settings.

**KEYWORDS:** Trauma, Mental health nursing, Inpatient psychiatry, Trauma-Focused Cognitive Behavioral Therapy, TF-CBT, Trauma-informed care, Psychiatric nursing, Post-traumatic stress disorder, Mental health treatment

## INTRODUCTION

In the field of psychiatric care, the impact of trauma on mental health has gained increasing attention in recent decades. Research consistently demonstrates that traumatic experiences, whether occurring in childhood or adulthood, can significantly influence the onset and progression of mental health disorders. For patients in inpatient psychiatric settings, trauma is often a contributing factor to the complex conditions they present with, including PTSD, depression, anxiety, and substance use disorders. Given the prevalence of trauma-related conditions, the integration of trauma-specific treatment modalities, such as Trauma-Focused Cognitive Behavioral Therapy (TF-CBT), is crucial to the recovery process.

Mental health nurses, as frontline healthcare providers, are uniquely positioned to play a critical role in the implementation of TF-CBT in inpatient psychiatric settings. Their frequent interactions with patients, their role in the ongoing management of care, and their ability to develop strong therapeutic relationships make them essential contributors to trauma-informed care. This article explores the significance of TF-CBT as a treatment for trauma, delves into the specific role of mental health nurses in administering this therapeutic approach, and examines the challenges and opportunities involved in implementing TF-CBT within inpatient psychiatric facilities.

### The Prevalence of Trauma in Psychiatric Inpatients

The prevalence of trauma in psychiatric inpatient populations is staggering. Studies estimate that between 75% to 90% of individuals receiving inpatient psychiatric care have experienced significant trauma, which often underlies their mental health disorders (Substance Abuse and Mental Health

Services Administration [SAMHSA], 2014). Trauma may manifest in various forms, including childhood abuse, domestic violence, sexual assault, accidents, or natural disasters. For many patients, these traumatic experiences serve as the catalyst for developing conditions such as PTSD, complex trauma, or other related disorders.

In psychiatric inpatient settings, trauma frequently complicates the course of treatment. Symptoms of trauma, such as hypervigilance, dissociation, flashbacks, and emotional numbing, can exacerbate other mental health conditions and make it difficult for patients to engage in treatment. Trauma can also influence patients' responses to treatment, with some individuals exhibiting heightened sensitivity to the environment, medical procedures, or authority figures. Consequently, the adoption of a trauma-informed approach, including the use of TF-CBT, is crucial for addressing the needs of these patients and fostering their recovery.

### Trauma-Informed Care

Trauma-informed care is a framework that recognizes the widespread impact of trauma and emphasizes the importance of creating a safe, supportive environment for individuals who have experienced trauma. This approach acknowledges that trauma can affect every aspect of a person's life, including their physical, emotional, and psychological well-being. It also highlights the need for healthcare providers, including mental health nurses, to understand the impact of trauma and to tailor their care accordingly.

The principles of trauma-informed care include safety, trustworthiness, collaboration, empowerment, and choice. For mental health nurses working in inpatient psychiatric settings,



these principles guide their interactions with patients and shape the delivery of TF-CBT. By adopting a trauma-informed approach, nurses can help to mitigate the retraumatization that can occur in psychiatric settings and support patients in their healing process.

### Trauma-Focused Cognitive Behavioral Therapy (TF-CBT)

TF-CBT is a structured, evidence-based treatment model designed to help individuals process traumatic memories and develop healthier coping mechanisms for managing trauma-related symptoms. Originally developed for children and adolescents, TF-CBT has been adapted for use with adults in both inpatient and outpatient psychiatric settings. The therapy typically involves 12 to 16 sessions, although the duration and structure can be modified based on the patient's needs and the context in which the therapy is delivered.

### Core Components of TF-CBT

TF-CBT combines traditional cognitive behavioral therapy (CBT) techniques with trauma-sensitive interventions. The primary components of TF-CBT include:

- 1. Psychoeducation:** Educating the patient and their family about the nature of trauma and its impact on mental health. This includes normalizing the patient's reactions to trauma and helping them understand that their symptoms are a natural response to abnormal experiences.
- 2. Relaxation Techniques:** Teaching patients relaxation and stress management skills, such as deep breathing, progressive muscle relaxation, and mindfulness. These techniques help patients regulate their physiological responses to trauma-related stress.
- 3. Affective Modulation:** Assisting patients in recognizing and managing their emotions, particularly those triggered by trauma. This involves helping patients identify, label, and express their emotions in a healthy way.
- 4. Cognitive Processing:** Encouraging patients to challenge and reframe distorted thoughts related to the trauma. This component focuses on identifying unhelpful beliefs (e.g., self-blame, guilt) and replacing them with more balanced, realistic cognitions.
- 5. Trauma Narrative:** One of the most critical aspects of TF-CBT is the development of a trauma narrative, in which the patient gradually recounts their traumatic experience in a controlled and supportive environment. This process helps to desensitize the patient to traumatic memories and facilitates the integration of those memories into their broader life story.
- 6. In Vivo Exposure:** When appropriate, patients may engage in in vivo exposure, which involves confronting trauma-related triggers in real life (e.g., revisiting places associated with the trauma) in a controlled and therapeutic manner.
- 7. Consolidation and Termination:** The final stage of TF-CBT involves consolidating the gains made during therapy and preparing the patient for the end of the therapeutic relationship. This phase includes relapse prevention strategies and a focus on future goals.

### The Role of Mental Health Nurses in Implementing TF-CBT

Mental health nurses are ideally positioned to play a key role in the implementation of TF-CBT in inpatient psychiatric settings. Their ongoing, close relationships with patients enable them to provide the consistent support required for effective trauma treatment. Furthermore, nurses are often responsible for delivering elements of trauma-informed care, even if they are not directly providing the full course of TF-CBT. Below, we examine the specific contributions of mental health nurses in delivering TF-CBT and enhancing trauma care.

#### 1. Screening for Trauma

The first step in the successful implementation of TF-CBT is the identification of patients who have experienced trauma. Mental health nurses are often the first point of contact for patients entering inpatient psychiatric care, and they play a vital role in conducting trauma screenings. These screenings typically involve the use of validated tools, such as the Trauma History Questionnaire (THQ) or the PTSD Checklist for DSM-5 (PCL-5). Early identification of trauma enables the care team to determine whether TF-CBT is an appropriate intervention for the patient.

During the screening process, nurses must create a safe and supportive environment in which patients feel comfortable disclosing their trauma histories. This requires nurses to be sensitive to the patient's emotional state and to use trauma-informed communication techniques, such as active listening, empathy, and non-judgmental responses. By establishing a foundation of trust, nurses can encourage patients to share their experiences and begin the process of healing.

#### 2. Providing Psychoeducation

One of the core components of TF-CBT is psychoeducation, which involves teaching patients and their families about the effects of trauma on mental health and the process of trauma-focused treatment. Mental health nurses are often responsible for delivering this information to patients and their families, particularly in the early stages of treatment.

Psychoeducation serves several important purposes. First, it helps to normalize the patient's symptoms and reduce feelings of shame or guilt. Many trauma survivors blame themselves for their experiences or feel as though they are "crazy" for having trauma-related symptoms. By providing accurate, compassionate information about the nature of trauma, nurses can help to alleviate these feelings and foster a sense of hope and empowerment.

Second, psychoeducation prepares patients and families for the TF-CBT process. Trauma treatment can be emotionally challenging, and it is important for patients to understand what to expect from therapy. Nurses play a key role in setting realistic expectations and addressing any concerns or fears that patients or their families may have about TF-CBT.

#### 3. Collaborating with the Multidisciplinary Team

In inpatient psychiatric settings, TF-CBT is often delivered as part of a multidisciplinary approach to care. Mental health nurses collaborate with psychologists, psychiatrists, social



workers, and other healthcare professionals to ensure that TF-CBT is integrated into the patient's overall treatment plan.

Nurses act as advocates for the patient, ensuring that trauma-informed care principles are upheld throughout the patient's stay. They also provide valuable insights into the patient's progress, symptoms, and responses to treatment. For example, if a patient is experiencing heightened anxiety during the trauma narrative phase of TF-CBT, the nurse can communicate this to the therapist and suggest modifications to the treatment plan. This collaborative approach ensures that TF-CBT is delivered in a way that meets the patient's individual needs and promotes healing.

#### 4. Building Therapeutic Relationships

Therapeutic rapport is essential for the success of TF-CBT and trauma-informed care in general. Mental health nurses spend more time with patients than any other member of the healthcare team, making them the primary source of emotional support for patients during their inpatient stay.

Building a therapeutic relationship requires nurses to create a safe and trusting environment in which patients feel comfortable discussing their trauma. This involves using trauma-informed communication techniques, such as active listening, validation, and empathy. Nurses must also be attuned to the patient's emotional state and respond to signs of distress with compassion and care.

By fostering a strong therapeutic relationship, nurses can help patients feel supported and understood, which is critical for engaging them in the TF-CBT process. Patients who feel safe and valued are more likely to participate fully in therapy and to make meaningful progress toward healing.

#### 5. Supporting Coping Strategies

Another key component of TF-CBT is the development of coping strategies to manage trauma-related symptoms, such as anxiety, depression, and emotional dysregulation. Mental health nurses work closely with patients to reinforce the coping techniques taught during TF-CBT sessions.

For example, nurses may guide patients through relaxation exercises, such as deep breathing or progressive muscle relaxation, during moments of heightened anxiety. They may also help patients identify triggers for trauma-related symptoms and develop strategies for managing those triggers in real-time. By providing ongoing support and reinforcement, nurses play a critical role in helping patients internalize and apply the coping strategies learned in TF-CBT.

In addition to providing direct support to patients, nurses also play a role in educating patients' families about coping strategies. Family involvement is often a key component of TF-CBT, and nurses can help families understand how to support their loved one's recovery process. This may include teaching family members how to provide emotional support, encouraging positive coping mechanisms, and fostering open communication about trauma.

#### 6. Monitoring Progress

Monitoring patient progress is an essential aspect of TF-CBT, as it allows the treatment team to assess the effectiveness of therapy and make adjustments as needed. Mental health nurses are responsible for tracking patients' responses to treatment and documenting any changes in symptoms, behaviors, or emotional states.

Nurses may use a variety of assessment tools to measure progress, including symptom checklists, mood scales, and behavioral observations. They also engage in ongoing dialogue with patients to gauge how they are feeling and to identify any challenges they may be experiencing with TF-CBT. This feedback is shared with the multidisciplinary team, allowing for timely adjustments to the treatment plan.

For example, if a patient is struggling with the trauma narrative phase of TF-CBT, the nurse may suggest that the therapist slow down the process or incorporate additional relaxation techniques to help the patient feel more comfortable. By closely monitoring progress and communicating with the treatment team, nurses help to ensure that TF-CBT is tailored to the patient's needs and that treatment goals are being met.

#### 7. Advocacy and Trauma-Informed Care

Mental health nurses are strong advocates for the integration of trauma-informed care principles in inpatient psychiatric settings. This includes promoting a safe and supportive environment for all patients, particularly those with trauma histories.

Nurses advocate for the reduction of coercive practices, such as restraints or seclusion, which can be retraumatizing for individuals with trauma histories. They also work to ensure that all staff members are aware of the potential impact of trauma on patient behavior and that they respond to patients with empathy and understanding.

In addition to advocating for individual patients, nurses also play a role in shaping institutional policies and practices related to trauma-informed care. For example, they may participate in staff training programs on trauma-informed care or contribute to the development of trauma-sensitive protocols for patient interactions and interventions.

By advocating for trauma-informed care at both the individual and institutional levels, mental health nurses help to create an environment in which TF-CBT can be delivered effectively and in which all patients feel safe and supported.

#### Barriers to Implementing TF-CBT in Inpatient Settings

While TF-CBT is a highly effective treatment for trauma, its implementation in inpatient psychiatric settings can be challenging. Several barriers can impede the successful delivery of TF-CBT, including time constraints, staff training needs, patient engagement challenges, and systemic issues. Addressing these barriers is essential for ensuring that patients in inpatient settings receive the trauma care they need.



## 1. Time Constraints

Inpatient psychiatric stays are often short, particularly in acute care settings. The average length of stay in an inpatient psychiatric facility is approximately 7 to 10 days, which may not provide enough time to complete a full course of TF-CBT. Given that TF-CBT typically involves 12 to 16 sessions, nurses and other healthcare providers may need to modify the therapy to fit within the patient's limited time in the facility.

One potential solution to this challenge is the development of a condensed version of TF-CBT that focuses on the most critical elements of the therapy, such as psychoeducation, relaxation techniques, and cognitive processing of trauma. Nurses can play a key role in delivering these components during the patient's inpatient stay and in coordinating follow-up care with outpatient providers to ensure that the patient continues TF-CBT after discharge.

## 2. Staff Training

TF-CBT is a specialized form of therapy that requires specific training and expertise. Not all mental health nurses may have received formal training in TF-CBT, which can limit their ability to deliver the therapy effectively. Furthermore, inpatient psychiatric facilities may not have the resources to provide ongoing professional development in trauma-focused interventions.

To address this barrier, inpatient facilities should invest in training programs that equip nurses with the knowledge and skills needed to deliver TF-CBT. This may include workshops, online courses, or mentorship opportunities with experienced TF-CBT providers. Additionally, facilities should encourage a culture of continuous learning, in which nurses are supported in their efforts to expand their expertise in trauma care.

## 3. Patient Engagement

Patients in inpatient psychiatric settings often present with severe mental health conditions, such as psychosis, severe depression, or suicidality, which can make it difficult for them to engage in structured therapy like TF-CBT. Some patients may struggle to concentrate during therapy sessions or may be emotionally overwhelmed by the process of revisiting their trauma.

Mental health nurses must use their clinical judgment to determine whether TF-CBT is appropriate for each patient and to identify any modifications that may be needed to accommodate the patient's current mental state. For example, nurses may need to break TF-CBT sessions into shorter, more manageable segments or incorporate additional coping strategies to help patients manage their emotional responses.

In cases where TF-CBT is not feasible, nurses can explore alternative trauma treatments, such as Eye Movement Desensitization and Reprocessing (EMDR) or Dialectical Behavior Therapy (DBT), which may be more appropriate for certain patients.

## 4. Systemic Issues

In some inpatient psychiatric settings, systemic issues, such as high patient-to-staff ratios, limited resources, and

organizational barriers, can impede the delivery of TF-CBT. Nurses may have limited time to spend with each patient, and there may be insufficient staff to provide the level of care required for trauma-focused interventions.

To overcome these challenges, inpatient facilities should prioritize trauma care and allocate the necessary resources to support its implementation. This may include hiring additional staff, providing trauma-specific training for all healthcare providers, and incorporating trauma-informed care principles into the facility's policies and procedures.

## Improving Patient Outcomes through Trauma-Informed Care

The implementation of TF-CBT by mental health nurses in inpatient psychiatric settings can significantly improve patient outcomes. Research has shown that trauma-informed care, including TF-CBT, leads to reductions in PTSD symptoms, decreased use of restraints and seclusion, and improved overall patient satisfaction (Green et al., 2016). By adopting TF-CBT as part of a comprehensive trauma-informed care strategy, nurses can provide more targeted and effective treatment for trauma-related disorders.

### 1. Reducing PTSD Symptoms

TF-CBT has been shown to be highly effective in reducing the symptoms of PTSD, including intrusive thoughts, flashbacks, hypervigilance, and emotional numbing. By helping patients process their traumatic experiences and develop healthier coping mechanisms, TF-CBT can alleviate the distress associated with trauma and promote recovery.

Mental health nurses play a key role in supporting patients throughout this process, providing the emotional support and guidance needed to navigate the challenges of trauma therapy. By fostering a safe and supportive environment, nurses can help patients feel more comfortable engaging in TF-CBT and making meaningful progress toward healing.

### 2. Decreasing the Use of Restraints and Seclusion

In inpatient psychiatric settings, the use of restraints and seclusion is often a last resort for managing patients who are experiencing extreme agitation or aggression. However, these practices can be retraumatizing for individuals with a history of trauma, exacerbating their symptoms and prolonging their stay in the facility.

Trauma-informed care, including TF-CBT, emphasizes the importance of minimizing the use of coercive practices and providing patients with alternative strategies for managing their distress. By teaching patients relaxation techniques, emotion regulation skills, and other coping mechanisms, nurses can help to reduce the need for restraints and seclusion.

### 3. Improving Patient Satisfaction

Patients who receive trauma-informed care, including TF-CBT, report higher levels of satisfaction with their treatment. They feel more understood, supported, and empowered to take an active role in their recovery. This sense of empowerment is particularly important for trauma survivors, who may have





experienced a loss of control or agency during their traumatic experiences.

By providing trauma-informed care and implementing TF-CBT, mental health nurses can help to create a more positive and healing environment for patients. This, in turn, can lead to better treatment outcomes and a more successful recovery process.

## CONCLUSION

Mental health nurses play an essential role in enhancing trauma care within inpatient psychiatric settings through the implementation of TF-CBT. Their ability to assess, educate, and collaborate with patients and multidisciplinary teams makes them integral to the trauma-informed care framework. By addressing the unique needs of trauma survivors through evidence-based approaches like TF-CBT, nurses can help improve patient outcomes, foster resilience, and contribute to long-term recovery. Despite the barriers, such as time constraints and the need for specialized training, the integration of TF-CBT into nursing practice offers a powerful tool for healing the deep wounds caused by trauma. Moving forward, healthcare institutions must invest in nurse education, support multidisciplinary collaboration, and address systemic barriers to ensure the widespread implementation of this vital therapeutic approach. In doing so, mental health nurses will continue to serve as advocates for trauma survivors, facilitating their journey toward recovery and well-being.

## BIBLIOGRAPHY

1. American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders (5th ed.)*. American Psychiatric Association.
2. Cohen, J. A., Mannarino, A. P., & Deblinger, E. (2016). *Trauma-focused cognitive-behavioral therapy for children and adolescents: Treatment applications*. Guilford Press.
3. de Arellano, M. A. R., Lyman, D. R., Jobe-Shields, L., George, P., Dougherty, R. H., Daniels, A. S., ... & Delphin-Rittmon, M. E. (2014). Trauma-focused cognitive-behavioral therapy for children and adolescents: Assessing the evidence. *Psychiatric services, 65*(5), 591-602.
4. Green, B. L., Kaltman, S., Frank, L., et al. (2016). Trauma-focused cognitive behavioral therapy for traumatized adults: A review of the evidence. *Behavior Therapy, 47*(5), 652-668. <https://doi.org/10.1016/j.beth.2015.05.005>
5. Substance Abuse and Mental Health Services Administration (SAMHSA). (2014). *SAMHSA's concept of trauma and guidance for a trauma-informed approach (HHS Publication No. SMA-14-4884)*. SAMHSA.
6. Foa, E. B., Keane, T. M., Friedman, M. J., & Cohen, J. A. (Eds.). (2008). *Effective treatments for PTSD: Practice guidelines from the International Society for Traumatic Stress Studies (2nd ed.)*. Guilford Press.
7. Ford, J. D., & Courtois, C. A. (2013). *Treating complex traumatic stress disorders in adults: Scientific foundations and therapeutic models*. Guilford Press.
8. Deblinger, E., Mannarino, A. P., Cohen, J. A., Runyon, M. K., & Steer, R. A. (2011). Trauma-focused cognitive behavioral therapy for children: impact of the trauma narrative and treatment length. *Depression and anxiety, 28*(1), 67-75.
9. Herman, J. L. (2015). *Trauma and recovery: The aftermath of violence--from domestic abuse to political terror*. Basic Books.
10. van der Kolk, B. A. (2014). *The body keeps the score: Brain, mind, and body in the healing of trauma*. Viking.
11. Deblinger, E., Pollio, E., & Dorsey, S. (2016). Applying trauma-focused cognitive-behavioral therapy in group settings. *Child Maltreatment, 21*(1), 59-73. <https://doi.org/10.1177/1077559515620668>
12. National Institute for Health and Care Excellence (NICE). (2018). *Post-traumatic stress disorder: Management*. NICE Guideline [NG116]. <https://www.nice.org.uk/guidance/ng116>
13. Cohen, J. A., Mannarino, A. P., Berliner, L., & Deblinger, E. (2000). Trauma-focused cognitive behavioral therapy for children and adolescents: An empirical update. *Journal of interpersonal violence, 15*(11), 1202-1223.
14. Courtois, C. A., & Ford, J. D. (2013). *Treatment of complex trauma: A sequenced, relationship-based approach*. Guilford Press.
15. Webb, C., Hayes, A. M., Grasso, D., Laurenceau, J. P., & Deblinger, E. (2014). Trauma-focused cognitive behavioral therapy for youth: Effectiveness in a community setting. *Psychological trauma: theory, research, practice, and policy, 6*(5), 555.
16. Beck, A. T. (2019). *Cognitive therapy of depression*. Guilford Press.
17. Shapiro, F. (2018). *Eye movement desensitization and reprocessing (EMDR) therapy: Basic principles, protocols, and procedures (3rd ed.)*. Guilford Press.
18. Cloitre, M., Courtois, C. A., Charuvastra, A., et al. (2012). Treatment of complex PTSD: Results of the ISTSS expert clinician survey on best practices. *Journal of Traumatic Stress, 24*(6), 615-627. <https://doi.org/10.1002/jts.20697>
19. Porges, S. W. (2017). *The pocket guide to the polyvagal theory: The transformative power of feeling safe*. Norton.
20. Briere, J. N., & Scott, C. (2015). *Principles of trauma therapy: A guide to symptoms, evaluation, and treatment (2nd ed.)*. Sage.
21. Jensen, T. K., Holt, T., Ormhaug, S. M., Egeland, K., Granly, L., Hoaas, L. C., ... & Wentzel-Larsen, T. (2014). A randomized effectiveness study comparing trauma-focused cognitive behavioral therapy with therapy as usual for youth. *Journal of Clinical Child & Adolescent Psychology, 43*(3), 356-369.
22. Zaleski, K. L., Johnson, D. M., & Klein, C. D. (2016). Exploring the impact of trauma-focused cognitive-behavioral therapy on emotional regulation in a military population. *Journal of Traumatic Stress, 29*(4), 297-303. <https://doi.org/10.1002/jts.22116>
23. National Center for PTSD. (2015). *Trauma-focused psychotherapy: Best practice for PTSD*. U.S. Department of Veterans Affairs. [https://www.ptsd.va.gov/professional/treat/txessentials/trauma\\_focused\\_tx.asp](https://www.ptsd.va.gov/professional/treat/txessentials/trauma_focused_tx.asp)
24. Kar, N. (2011). Cognitive behavioral therapy for the treatment of post-traumatic stress disorder: A review. *Neuropsychiatric Disease and Treatment, 7*, 167-181. <https://doi.org/10.2147/NDT.S10389>
25. Ford, J. D., Grasso, D. J., Greene, C. A., et al. (2018). Evidence-based practices for trauma and traumatic stress in children and adolescents. Springer Publishing.
26. Scheeringa, M. S., Weems, C. F., Cohen, J. A., Amaya-Jackson, L., & Guthrie, D. (2011). Trauma-focused cognitive-behavioral therapy for posttraumatic stress disorder in three-through six year-old children: A randomized clinical trial. *Journal of Child Psychology and Psychiatry, 52*(8), 853-860.